

Cultural Information for:	Anemone Gemstone	Annual
Common Name:	Adonis or Windflower	
Botanical Name:	Anemone coronaria	
Seed Count:	45,000-57,000/ounce	1,600-2,000/gram
Optimum Germination Temperature:	59°F / 15°C (Chamber)	
Optimum Growing Temperature:	50-75°F / 10-24°C	
Optimum pH:	5.8 – 6.2 (plug/peat moss)	6.0 – 7.0 (finishing/mineral soil)
EC – Plug:	0.26 – 0.75 mmhos/cm (1:2) / 0.76 – 2.0 (SME)	
EC – Finishing:	0.76 – 1.0 mmhos/cm (1:2) / 2.1 – 3.0 (SME)	

Plug Culture: 8 weeks (288 / 12 x 24 deep tray)

Stage One (days 1 – 14) Sow seed into a 288 deep plug tray using a well-drained sterile media with good aeration. Place uncovered in a germination chamber and drench with 100 ppm's of Captan to avoid disease (damping off and botrytis). Optimum germination temperature is 59°F/15°C. Provide high humidity throughout the germination period, never allowing the media to dry out. After radicle emergence (6-7 days), lightly cover with a peat/vermiculite blend and provide light in the chamber. If germinating in a greenhouse lightly cover with a peat/vermiculite blend at sowing and avoid temperatures above 65°F/18°C.

Stage Two (day 15-35) The cotyledons are fully expanded. Place the plugs in a well-ventilated greenhouse with low humidity to avoid disease problems. Provide a light level of 2,000-2,500 foot candles/22,000-27,000 lux. Optimum growing temperature is 60-65°F / 16-18°C at night and less than 75°F / 24°C during the day. Fertilize 2-3 times per week at 50-75 ppm N using a well-balanced formulation such as 15-16-17.

Stage Three (day 35-49) As the plugs begin to fill in the trays, allow the plants to dry down slightly in between irrigations to maximize root growth. Increase fertilizer rate to 100- 150 ppm of N as needed to maintain healthy growth.

Stage Four (days 50-56) Increase light level to 2,500-5,000 f.c./ 27,000-54,000 lux to acclimate prior to transplanting into cut flower beds. Applying extra phosphorus at this stage will strengthen the root system prior to transplanting.

Finishing: 12-14 weeks

Transplanting: When the plugs have 4-6 true leaves, they are ready to transplant into cut flower beds. Take special care in removing the plugs from the trays to avoid root damage. Anemones will respond best in a soil with good fertility, high organic matter and excellent drainage. Optimum pH in soil-based media is 6.0-7.0.

Spacing: Recommended spacing is 6" x 6" (37 plants per square yard) / 15 cm x 15 cm. (44 plants per square meter).

Fertilizer: Maintain the soil somewhat dry at first to promote new root growth. Once new roots emerge, apply 150- 200 ppm N from a well-balanced liquid fertilizer, such as 15-16-17, to promote healthy growth of the plants. Periodic applications of calcium nitrate are recommended to strengthen flower stems and plant tissue. Water thoroughly to prevent excess salts. Excess nitrogen can promote overgrowth of the foliage. Water early in the day if irrigating overhead.

Temperature: After transplanting, establish at 60-65°F / 15-18°C for one week. Next, lower the temperature to 50-55°F / 10-13°C nighttime and 59-65°F / 15-18°C daytime for optimum development and flower stem length. Ideally, do not exceed 68°F / 20°C daytime and below 41°F / 5°C nighttime.

Light: Anemone grow best at 3,500-5,000 f.c. / 38,000-54,000 lux. Apply shading as needed to reduce excess heat from intense sunshine or to increase stem length in high light areas. Anemone Gemstone is day length neutral but develops faster under longer photoperiods and warmer temperatures.

Insect: Aphids, thrips and whiteflies

Disease: Botrytis, colletotrichum (leaf curl), downy mildew, pythium and rhizoctonia. Overhead water before noon to allow the foliage to dry before evening.

Crop Schedule: Sow from July to September for flowering from January to June. Total crop time from sowing to first flowering is 20-22 weeks. Year-round sowing in cool equatorial regions.

Harvesting: Flowers should be harvested just prior to opening. Harvest when the petals begin to separate from the center of the flower but not all the way open.

Post-Harvest: Keep stems in a 2-4% sugar solution or commercial holding solution to extend vase life. Recut stem ends during each transfer to extend the vase life. Keep flowers wrapped during rehydration for straighter stems. Store cut stems dry at 32-34°F / 0-1°C for a week. After conditioning, store stems in a holding solution containing sugar and a germicide to improve flower opening. Stems should be stored and shipped in a vertical position. Sensitive to ethylene.

"A true jewel among anemones: Gemstone produces over 20 flowers per plant, flowering under low light levels with no need for netting or disbudding".

"All information given is intended for general guidance only and may have to be adjusted to meet individual needs. Cultural details are based on North American conditions and Sakata cannot be held responsible for any crop damage related to the information given herein. Application of recommended growth regulators and chemicals are subject to local and state regulations. Always follow manufacturer's label instructions. Testing a few plants prior to treating the entire crop is best."